

## OBSTETRICS

# Enoxaparin for the prevention of preeclampsia and intrauterine growth restriction in women with a history: a randomized trial



Katie M. Groom, MBBS, PhD, FRANZCOG, CMFM; Lesley M. McCowan, MB ChB, MD, FRANZCOG, CMFM; Laura K. Mackay, BSc; Arier C. Lee, PhD; Joanne M. Said, MBBS, PhD, FRANZCOG, CMFM; Stefan C. Kane, MBBS, FRANZCOG; Susan P. Walker, MBBS, MD, FRANZCOG, CMFM; Thijs E. van Mens, MD, MBA; Natalie J. Hannan, PhD; Stephen Tong, MBBS, PhD, FRANZCOG; Larry W. Chamley, PhD; Peter R. Stone, MB ChB, MD, FRANZCOG, CMFM; Claire McLintock, MB ChB, FRACP; the Enoxaparin for Prevention of Preeclampsia and Intrauterine Growth Restriction Trial Investigator Group



**BACKGROUND:** Preeclampsia and small-for-gestational-age pregnancy are major causes of maternal and perinatal morbidity and mortality. Women with a previous pregnancy affected by these conditions are at an increased risk of recurrence in a future pregnancy. Past trials evaluating the effect of low-molecular-weight heparin for the prevention of recurrence of preeclampsia and small-for-gestational-age pregnancy have shown conflicting results with high levels of heterogeneity displayed when trials were compared.

**OBJECTIVE:** We sought to assess the effectiveness of enoxaparin in addition to high-risk care for the prevention of preeclampsia and small-for-gestational-age pregnancy in women with a history of these conditions.

**STUDY DESIGN:** This was an open-label randomized controlled trial in 5 tertiary care centers in 3 countries. Women with a viable singleton pregnancy were invited to participate between  $>6^{+0}$  and  $<16^{+0}$  weeks if deemed to be at high risk of preeclampsia and/or small for gestational age based on their obstetric history. Eligible participants were randomly assigned in a 1-to-1 ratio to standard high-risk care or standard high-risk care plus enoxaparin 40 mg (4000 IU) by subcutaneous injection daily from recruitment until  $36^{+0}$  weeks or delivery, whichever occurred sooner. Standard high-risk care was defined as care coordinated by a high-risk antenatal clinic service, aspirin 100 mg daily until  $36^{+0}$  weeks, and—for women with prior preeclampsia—calcium 1000–1500 mg daily until  $36^{+0}$  weeks. In a subgroup of participants serum samples were taken at recruitment and at 20 and 30 weeks' gestation and later analyzed for soluble fms-like tyrosine kinase-1, soluble endoglin, endothelin-1, placental growth factor, and soluble vascular cell adhesion molecule 1. The primary outcome was a composite of preeclampsia and/or

small-for-gestational-age  $<5$ th customized birthweight percentile. All data were analyzed on an intention-to-treat basis. The trial is registered with the Australian New Zealand Clinical Trials Registry (ACTRN12609000699268).

**RESULTS:** Between July 26, 2010, and Oct. 28, 2015, a total of 156 participants were enrolled and included in the analysis. In all, 149 participants were included in the outcome analysis (72 receiving standard high-risk care plus enoxaparin and 77 receiving standard high-risk care only). Seven women who miscarried  $<16$  weeks' gestation were excluded. The majority of participants (151/156, 97%) received aspirin. The addition of enoxaparin had no effect on the rate of preeclampsia and/or small-for-gestational-age  $<5$ th customized birthweight percentile: enoxaparin 18/72 (25%) vs no enoxaparin 17/77 (22.1%) (odds ratio, 1.19; 95% confidence interval, 0.53–2.64). There was also no difference in any of the secondary outcome measures. Levels of soluble fms-like tyrosine kinase-1 and soluble endoglin increased among those who developed preeclampsia, but there was no difference in levels of these antiangiogenic factors (nor any of the other serum analytes measured) among those treated with enoxaparin compared to those receiving standard high-risk care only.

**CONCLUSION:** The use of enoxaparin in addition to standard high-risk care does not reduce the risk of recurrence of preeclampsia and small-for-gestational-age infants in a subsequent pregnancy.

**Key words:** enoxaparin, fetal growth restriction, intrauterine growth restriction, low-molecular-weight heparin, preeclampsia, randomized trial, small for gestational age

## Introduction

Preeclampsia and intrauterine growth restriction (IUGR) are common causes

## EDITORS' CHOICE

of maternal and perinatal morbidity and mortality. Preeclampsia complicates 3–5% of pregnancies. IUGR is more difficult to define and measure but approximately 10% of infants will be born small for gestational age (SGA) defined as birthweight  $<10$ th customized birthweight percentile. At least two thirds of these infants will have had evidence of abnormal uterine and umbilical artery Doppler waveforms if

diagnosed prior to birth suggesting significant uteroplacental disease.<sup>1</sup> Thus SGA is used as the most reliable surrogate marker for IUGR.

There are a wide variety of identified risk factors for preeclampsia and/or SGA but risk prediction models, at best, remain modest.<sup>2,3</sup> Until relatively recently it has been suggested that inherited thrombophilias are associated with preeclampsia and SGA, however, more recent evidence from prospective cohort studies suggests this association,

**Cite this article as:** Groom KM, McCowan LM, Mackay LK, et al. Enoxaparin for the prevention of preeclampsia and intrauterine growth restriction in women with a history: a randomized trial. *Am J Obstet Gynecol* 2017;216:296.e1–14.

0002-9378/free

© 2017 Elsevier Inc. All rights reserved.

<http://dx.doi.org/10.1016/j.ajog.2017.01.014>

if present, is only weak.<sup>4</sup> Obstetric history remains the most commonly used method for risk assessment in current clinical practice. Women with previous preeclampsia and/or SGA are at significant risk of recurrence especially when the disease was severe and occurred early in pregnancy.<sup>5-8</sup>

Preeclampsia and IUGR are considered placental diseases and there is likely to be considerable overlap in pathological mechanisms. This fact has led investigators to research common therapeutic and preventative strategies for both diseases. Aspirin and calcium have been studied in a large number of randomized trials in a variety of populations and although effect sizes are only modest both significantly reduce the incidence of preeclampsia,<sup>9,10</sup> and aspirin also decreases the incidence of SGA.<sup>11</sup> Their use should be considered standard practice for women at high risk of these conditions.<sup>12-14</sup>

Heparin and low-molecular-weight heparin (LMWH) have potential as preventative therapies. Their presumed benefit may relate to their anticoagulant properties although it is likely that additional effects on trophoblast development may be more important.<sup>15-17</sup> Small observational and non-randomized trials<sup>18-21</sup> reporting benefit have led to a variety of randomized controlled trials. These initially focused specifically on populations with or without thrombophilia,<sup>22-24</sup> but trials commenced more recently have included women regardless of thrombophilia status. Results from individual trials of LMWH are conflicting,<sup>22-28</sup> possibly reflecting the heterogeneity of the populations being examined, the type of heparin/LMWH being tested, prolonged trial recruitment phases,<sup>23,24</sup> and early trial discontinuation due to presumed overwhelming effect<sup>22</sup> or futility of ability show any effect.<sup>25</sup> Recent meta-analysis<sup>29</sup> and individual patient data meta-analysis<sup>30</sup> also failed to conclusively demonstrate that LMWH reduces the risk of placenta-mediated complications in subsequent pregnancies for those deemed to be at high risk.

The aim of the Enoxaparin for the Prevention of Preeclampsia and IUGR

(EPPI) trial was to assess the effectiveness of LMWH for the prevention of recurrence of preeclampsia and SGA. The trial aimed to be more precise, and clinically relevant, with its inclusion criteria and primary outcome measures specific to women at high risk of preeclampsia and/or SGA. Account was made of each participant's thrombophilia status but this status did not define the study population. All participants received high-risk care including the use of low-dose aspirin and, where appropriate, calcium.

The EPPI trial is the first randomized controlled trial of LMWH to report the serial assessment of placentally derived angiogenic growth factors involved in the pathophysiological process of preeclampsia (soluble fms-like tyrosine kinase [sFlt]-1 and soluble endoglin [sEng], which are elevated in preeclampsia<sup>31-34</sup>; placental growth factor [PlGF], which is decreased in preeclampsia) as well as endothelial-derived circulating markers that are associated with maternal endothelial dysfunction (endothelin [ET]-1 and soluble vascular cell adhesion molecule [sVCAM]-1).

## Materials and Methods

### Study design, setting, and ethics statement

This was a multicenter open-label randomized controlled trial (ACTRN12609000699268) at 5 tertiary care centers in New Zealand, Australia, and The Netherlands. Appropriate ethics and governance approvals were obtained at each center. All women participating in the trial provided written informed consent.

### Participants

Through the duration of the trial women referred for antenatal care were screened for eligibility. Women were eligible for inclusion if they were  $>6^{+0}$  and  $<16^{+0}$  weeks; gestation with a viable singleton pregnancy confirmed by ultrasound scan and at risk of preeclampsia and/or IUGR based on their obstetric history with: (1) previous preeclampsia delivered  $<36^{+0}$  weeks in their last ongoing pregnancy reaching  $>12$  weeks; or (2) previous SGA infant  $<10$ th customized birthweight percentile delivered  $<36^{+0}$  weeks

in their last ongoing pregnancy reaching  $>12$  weeks with no major fetal anomaly; or (3) previous SGA infant  $\leq 3$ rd customized birthweight percentile delivered at any gestation in their last ongoing pregnancy reaching  $>12$  weeks with no major fetal anomaly. Eligibility criteria were checked against medical records. Women were excluded from the trial if they met  $\geq 1$  of the exclusion criteria: any contraindication to LMWH use; need for anticoagulant use in pregnancy such as previous thrombosis or antiphospholipid syndrome; previous successful pregnancy with LMWH treatment; multiple pregnancy; known preexisting type 1 or 2 diabetes; renal disease (with serum creatinine  $>150$   $\mu\text{mol/L}$ ); thrombocytopenia (platelet count  $<80 \times 10^9/\text{L}$ ); or known major fetal anomaly/chromosomal abnormality.

We included women with previous preeclampsia and/or SGA as these diseases are both placental in origin with considerable overlap in pathological mechanisms. LMWH is likely to exert any therapeutic benefit via effect(s) on placentation and therefore has potential to impact both diseases, which often coexist, particularly when disease is preterm.

### Randomization and interventions

After confirming eligibility and obtaining consent participants were randomly assigned in a 1-to-1 ratio to standard high-risk care or standard high-risk care plus enoxaparin 40 mg (4000 IU) by subcutaneous injection (Clexane, Sanofi-Aventis, Auckland, New Zealand)<sup>35</sup> daily from recruitment until  $36^{+0}$  weeks or delivery, whichever occurred sooner. A 40-mg dose of enoxaparin was selected as the standard dose used for venous thromboembolism prophylaxis and, as per manufacturer's direction, no adjustment was made for body mass index. Standard high-risk care was defined as care coordinated by a high-risk antenatal clinic service, aspirin 100 mg daily until  $36^{+0}$  weeks, and—for women with prior preeclampsia—calcium 1000–1500 mg daily until  $36^{+0}$  weeks. This was an open-label trial with all participants, clinicians, and investigators aware of trial group assignment.

A computer-generated randomization program balanced in blocks of 5 was

used with stratification for recruiting site and thrombophilia status. Participants who were tested prior to trial involvement were assigned to: (1) positive thrombophilia or (2) negative thrombophilia. Women not tested or partially tested (with negative result) were assigned: (3) unknown thrombophilia status. Samples were then taken prior to any use of LMWH and tested for lupus anticoagulant, anticardiolipin antibodies, antithrombin III deficiency, protein C deficiency, protein S deficiency, factor V Leiden, and the prothrombin gene mutation. These results were used in the final analysis but not revealed to participant, clinicians, or investigators during the trial period.

All participants were assigned a sequential trial identifying number according to thrombophilia status. At the lead recruiting site (National Women's Health, Auckland City Hospital) randomization occurred by telephone to the hospital pharmacy clinical trials service and trial group allocation was made by study identifying number. In all other sites sequential sealed opaque envelopes for each study identifying number were opened to reveal treatment allocation.

## Procedures

For women receiving LMWH, an educational session regarding injection technique was provided by a research midwife prior to treatment commencement. LMWH was resupplied on a monthly basis from hospital pharmacies. Participants were asked to return sharps disposal boxes with used syringes and any unused treatment. Indications to stop treatment  $<36^{+0}$  weeks included: need for delivery  $<36^{+0}$  weeks (stopped once decision made for delivery or 12 hours prior to induction of labor or elective cesarean delivery, whichever was later), episode of threatened preterm labor (treatment recommenced if symptoms settled), and clinical evidence of placental abruption or thrombocytopenia with platelet count  $<80 \times 10^9/L$ .

Serum samples were taken at recruitment (used as baseline) and at 20 and 30 weeks' gestation in a subgroup of participants ( $n = 127$ ). Samples were centrifuged and stored at  $-80^\circ C$  for later

assessment of sFlt-1, sEng, PlGF, sVCAM-1, and ET-1 by enzyme-linked immunosorbent assay according to manufacturer's instructions (R&D Systems, Minneapolis, MN). Optical density was determined using X-Mark microplate spectrophotometer (BioRad, Gladesville, Australia) and analyte levels were determined using Microplate manager 6 software (BioRad). Analytes were measured in 1 batch in a blinded fashion with the cases and controls mixed.

Standard fetal growth parameters<sup>36,37</sup> and uterine and umbilical arterial Doppler waveforms<sup>38</sup> were recorded at 20 and 24 weeks' gestation. Data from any additional antenatal ultrasound scans were collected. All participants received care through a high-risk antenatal clinic service including daily aspirin and, for women with prior preeclampsia, daily calcium. Care included regular blood pressure (BP) assessment, urine dipstick analysis for proteinuria, fetal well-being assessment, and review of any adverse events. Intrapartum and postnatal care was provided by local clinicians. Pregnancy, labor, postnatal, and neonatal data were collected by research staff from maternal and infant clinical records up to the time of hospital discharge.

## Outcomes

The primary outcome was a composite of preeclampsia and/or SGA  $<5$ th percentile. All cases with primary outcome events were reviewed by trial investigators. Secondary outcomes included preeclampsia, severe preeclampsia, HELLP syndrome (hemolysis, elevated liver enzyme, and low platelet count), eclampsia, gestational hypertension, SGA  $<10$ th percentile, SGA  $<5$ th percentile, SGA  $<3$ rd percentile, placental abruption and antepartum hemorrhage, thrombocytopenia (platelet count  $<100 \times 10^9/L$ ) while on LMWH, stillbirth, induction of labor, cesarean delivery, postpartum hemorrhage, gestational age at birth, preterm birth, mean birthweight and mean birthweight percentile, neonatal death, neonatal intensive care admission and duration of admission, a composite outcome of severe neonatal morbidity (evidence of  $\geq 1$ :

intraventricular hemorrhage [grade 3 and 4], cystic periventricular leukomalacia, chronic lung disease, retinopathy of prematurity requiring treatment, or necrotizing enterocolitis requiring surgery), and maternal serum levels of sFlt-1, sEng, PlGF, sVCAM-1, and ET-1.

Preeclampsia was defined as new-onset hypertension (systolic BP  $\geq 140$  mm Hg and/or diastolic BP  $\geq 90$  mm Hg)  $>20$  weeks' gestation with evidence of significant proteinuria or any multisystem complication including hematological, liver, renal, and neurological involvement.<sup>39</sup> Severe preeclampsia included preeclampsia associated with maternal death, persistent severe hypertension (systolic BP  $\geq 170$  mm Hg and/or diastolic BP  $\geq 110$  mm Hg), or a multisystem complication. Customized birthweight percentiles adjusting for infant sex, gestation at delivery, and maternal variables (parity, ethnicity, height, and early pregnancy weight) were used to identify SGA infants.<sup>40</sup> Nonpregnant- and pregnancy-specific (as appropriate to timing of testing) reference ranges were used to define protein C deficiency, protein S deficiency, and antithrombin III deficiency.<sup>41</sup>


## Statistical analysis

A priori we estimated a risk of recurrence of preeclampsia and/or SGA  $<5$ th percentile for the included population of 25%. Based on published data we estimated the risk of recurrence would be reduced to 7% with the addition of LMWH therapy.<sup>22</sup> We planned for a sample size of 160 participants (80 participants in each group, including a 5% dropout/early miscarriage rate) to achieve 80% power at a 2-sided significance level of .05 to detect a difference between 25-7%.

An independent data monitoring committee undertook a planned interim analysis once 98 participants were recruited and completed follow-up. Safety and primary outcome data were assessed and reported to the trial investigator group. Early discontinuation of the trial was to be considered if there was an excess of serious adverse events in the LMWH group or if a significant benefit had already been demonstrated (49

FIGURE 1

## Enoxaparin for Prevention of Preeclampsia and Intrauterine Growth Restriction (EPPI) trial profile



CONSORT flow diagram of participants in EPPI trial.

Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.

**TABLE 1**  
**Maternal and pregnancy characteristics**

|   | Standard high-risk care<br>and enoxaparin, n = 73 | Standard high-risk<br>care only, n = 83 |
|---|---|---|
| Baseline maternal characteristics                 |   |   |
| Maternal age, y                                   | 34 (21–42)  | 34 (20–40)                              |
| Ethnicity   |   |   |
| European/Caucasian                                | 50 (68.5%)  | 50 (60.2%)                              |
| Far East/South East Asian                         | 7 (9.6%)  | 5 (6.0%)                                |
| Maori/Cook Island                                 | 4 (5.5%)  | 4 (4.8%)                                |
| Pacific Island                                    | 1 (1.4%)  | 3 (3.6%)                                |
| South Asian                                       | 7 (9.6%)  | 7 (8.3%)                                |
| Other   | 4 (5.5%)  | 14 (6.9%)                               |
| Body mass index, kg/m <sup>2</sup>                | 26.3 (19–48)                                      | 25.1 (16–43)                            |
| Smoking status at trial entry                     | 3 (4.1%)  | 4 (4.8%)                                |
| Preexisting hypertension                          | 9 (12.3%)   | 9 (10.8%)                               |
| Previous gestational diabetes <sup>a</sup>        | 1 (1.4%)  | 3 (3.6%)                                |
| Blood pressure at trial entry                     |   |   |
| Systolic, mm Hg                                   | 110 (80–156)                                      | 110 (84–148)                            |
| Diastolic, mm Hg                                  | 70 (48–98)  | 69 (40–88)                              |
| Thrombophilia status                              |   |   |
| FVL heterozygous                                  | 6 (7.0%)  | 7 (8.4%)                                |
| Prothrombin gene<br>mutation heterozygous         | 3 (4.1%)  | 2 (2.4%)                                |
| Protein C deficiency                              | 1 (1.4%)  | 1 (1.2%)                                |
| Protein S deficiency                              | 5 (6.9%)  | 6 (7.2%)                                |
| Antithrombin III deficiency                       | 0   | 1 (1.2%)                                |
| FVL homozygous                                    | 0   | 0                                       |
| Prothrombin gene<br>mutation homozygous           | 0   | 0                                       |
| Lupus anticoagulant/anticardiolipin<br>antibodies | 0   | 3 (3.6%)                                |
| Incomplete screen/unreported result               | 7 (9.6%)  | 9 (10.8%)                               |
| Obstetric history and reasons for trial inclusion |   |   |
| Gravidity   |   |   |
| 2   | 36 (49.3%)  | 43 (51.8%)                              |
| 3   | 21 (28.8%)  | 18 (21.7%)                              |
| 4–5   | 12 (16.4%)  | 16 (19.2%)                              |
| ≥6  | 4 (5.5%)  | 6 (7.2%)                                |
| Parity  |   |   |
| 0   | 0   | 1 (1.2%) <sup>c</sup>                   |
| 1   | 63 (86.3%)  | 63 (75.9%)                              |
| 2   | 6 (8.2%)  | 14 (16.9%)                              |
| ≥3  | 4 (5.5%)  | 5 (6.0%)                                |

Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.

(continued)

participants in each group to achieve 80% power at a 2-sided significance level of .05 to detect a difference between 25–4%).

All data were analyzed on an intention-to-treat basis. Continuous data are reported as median with minimum–maximum values (range). Categorical data are reported as number and proportion. Unless otherwise stated, statistical models were adjusted for thrombophilia status, recruitment center, inclusion criteria, and number of previous preeclampsia/SGA events (1 vs ≥2) as these variables were deemed the most likely to potentially influence the risk of recurrence and/or the care provided. Multiple logistic regression was used to analyze the binary primary and secondary outcomes. Multiple linear regression was used to analyze the continuous secondary outcomes. No missing data imputation was conducted. No multiple comparison adjustment was made. Two-sided *P* values <.05 were used to determine statistical significance and all confidence intervals (CI) are given at a 2-sided 95% level. Statistical analysis was conducted using software (SAS for Windows 9.4; SAS Institute Inc, Cary, NC). Levels of analytes measured in the serum samples were tested for normal distribution and statistically tested using nonparametric Mann-Whitney test. Data were expressed as mean ± SEM. Biomarker statistical analysis was performed using software (GraphPad Prism 6; GraphPad Software, La Jolla, CA).

## Results

From July 26, 2010, through Oct. 28, 2015, a total of 157 women in 160 pregnancies consented and were randomly allocated to standard high-risk care and enoxaparin or standard high-risk care only. Three women were approached in 2 pregnancies and only the first eligible pregnancy was included; a fourth woman was deemed ineligible after randomization and was also excluded (156 participants were included). The outcome analysis included all pregnancies that proceeded >16 weeks (n = 149) (Figure 1).

Baseline characteristics for groups are reported in Table 1. Of participants, 44%



**TABLE 1**  
**Maternal and pregnancy characteristics** (continued)

|  | Standard high-risk care<br>and enoxaparin, n = 73 | Standard high-risk<br>care only, n = 83 |
|--|---|---|
| ≥2 Previous pregnancies affected by preeclampsia and/or SGA                                | 5 (6.9%)  | 15 (18.1%)                              |
| Inclusion criteria for trial entry <sup>b</sup>  |   |   |
| Previous preeclampsia delivered <36 <sup>+0</sup> wk                                       | 30 (41.1%)  | 38 (45.8%)                              |
| Previous SGA infant <10th customized birthweight percentile delivered <36 <sup>+0</sup> wk | 48 (65.8%)  | 43 (51.8%)                              |
| Previous SGA infant ≤3rd customized birthweight percentile delivered at any gestation      | 60 (82.2%)  | 63 (75.9%)                              |
| Pregnancy characteristics  |   |   |
| Gestational age at randomization, d  | 83 (42–111)                                       | 84 (42–111)                             |
| Aspirin use in pregnancy   | 71 (97.3%)  | 80 (96.4%)                              |
| Calcium use in pregnancy   | 43 (58.9%)  | 40 (48.2%)                              |
| Pregnancy multivitamin including folic acid use in pregnancy                               | 72 (98.6%)  | 77 (92.3%)                              |
| Miscarriage <16 wk (excluded from outcome analysis)  | 1 (1.4%)  | 6 (7.3%)                                |

Data expressed as median (minimum–maximum values) or number (percentage).

FVL, factor V Leiden; SGA, small for gestational age.

<sup>a</sup> Type 1/2 diabetes excluded from trial; <sup>b</sup> Each participant may meet >1 criteria; <sup>c</sup> Eligibility pregnancy was termination of pregnancy <20 wk gestation for severe early-onset intrauterine growth restriction.Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.

(68/156) had a history of preeclampsia delivered <36<sup>+0</sup> weeks and 58% (91/156) had a history of delivering a SGA infant <36<sup>+0</sup> weeks; 79% (123/156) had a history of delivering a severe SGA infant (≤3rd percentile). The large majority of women (151/156, 97%) received aspirin therapy during their pregnancy and 53% (83/156) received calcium therapy.

For women in the standard high-risk care and enoxaparin group the median duration of enoxaparin treatment was 156 days (range 1–209 days). Ten women reported skin reactions or significant bruising with enoxaparin treatment; 1 woman had an allergic reaction and transferred to dalteparin therapy. Twelve women elected to discontinue treatment <36<sup>+0</sup> weeks' gestation (6 due to concerns with injections, 6 due to maternal request for other reasons). No women

had thrombocytopenia while on enoxaparin therapy.

Overall, 35 women (23.5%) experienced the primary outcome of preeclampsia and/or SGA <5th percentile. There was no significant difference in the rates of the primary outcome between the 2 groups: 25% (18/72) in standard high-risk care and enoxaparin group compared with 22.1% (17/77) in standard high-risk care only group (adjusted odds ratio, 1.19; 95% CI, 0.53–2.64) (Table 2). There was also no significant difference in any secondary pregnancy, delivery, or neonatal outcomes (Tables 2–4). In subgroup analyses the use of enoxaparin had no effect on the rates of the primary outcome for women with a history of preeclampsia, SGA <3rd percentile, or SGA <10th percentile (Table 5).

Levels of sFlt-1, sEng, PlGF, sVCAM-1, and ET-1 in serum collected from

women at recruitment, 20 weeks, and 30 weeks (samples from 127 participants) showed no significant difference with enoxaparin treatment when compared to those receiving standard high-risk care only (Figure 2). Importantly, we identified a significant increase in sFlt-1, sEng, and sVCAM-1 levels in the 30-week time-point serum of women who developed preeclampsia, compared to those who did not (Figure 3). Levels of ET-1 were significantly increased in women who developed preeclampsia at all time points assessed (Figure 3). Among only those women who subsequently developed preeclampsia, no significant difference was observed in analytes in those treated with enoxaparin vs those receiving standard high-risk care only (Supplementary Figure 1).

Analysis of serum from women with pregnancies that subsequently developed SGA <5th percentile demonstrated no significant difference in sFlt-1, sEng, PlGF, or sVCAM-1 (Supplementary Figure 2, A–D). Like for those who developed preeclampsia, there was a significant increase in ET-1 levels at recruitment in the serum from women who later developed SGA pregnancy (Supplementary Figure 2, E).

## Comment

The EPPI trial has not shown any beneficial effect of enoxaparin use in pregnancy to reduce the risk of recurrence of preeclampsia and SGA. There may be several reasons for this real or apparent lack of effect.

Many of the previous trials of LMWH have had very broad inclusion criteria for the prevention of placenta-mediated pregnancy complications including recurrent miscarriage,<sup>27,28</sup> second-trimester loss,<sup>22,25</sup> placental abruption,<sup>42</sup> and unexplained stillbirth and nonpregnancy-related conditions (venous thromboembolism, phlebitis, and family history of thromboembolism).<sup>24</sup> If LMWH was effective in one subgroup but not others the significance of this effect may be lost within the pooled results. For this reason our trial aimed to be more specific to preeclampsia and SGA using clearly defined inclusion criteria and a limited

**TABLE 2**  
**Pregnancy outcomes**

|  | Standard high-risk care and enoxaparin, n = 72 | Standard high-risk care only, n = 77 | Adjusted odds ratio <sup>a</sup> (95% CI) |
|--|--|--------------------------------------|---|
| Primary outcome (preeclampsia and/or SGA infant <5th percentile) | 18 (25.0%)                                     | 17 (22.1%)                           | 1.19 (0.53–2.64)                          |
| Preeclampsia   | 6 (8.3%)                                       | 5 (6.5%)                             | 1.24 (0.33–4.64)                          |
| Severe preeclampsia  | 2 (2.8%)                                       | 1 (1.3%)                             | <sup>b</sup>                              |
| Preeclampsia with delivery $\leq 33^{+6}$ wk                     | 2 (2.8%)                                       | 1 (1.3%)                             | <sup>b</sup>                              |
| Preeclampsia with delivery $\leq 36^{+6}$ wk                     | 4 (5.6%)                                       | 2 (2.6%)                             | <sup>b</sup>                              |
| SGA infant <10th percentile                                      | 23 (31.9%)                                     | 23 (29.9%)                           | 1.17 (0.56–2.47)                          |
| SGA infant <5th percentile                                       | 15 (20.8%)                                     | 13 (16.9%)                           | 1.48 (0.61–3.62)                          |
| SGA infant <3rd percentile                                       | 9 (12.5%)                                      | 10 (13.0%)                           | 1.19 (0.40–3.52)                          |
| HELLP syndrome   | 1 (1.4%)                                       | 2 (2.6%)                             | <sup>b</sup>                              |
| Eclampsia  | 0  | 0                                    | NA  |
| Gestational hypertension   | 10 (14.9%)                                     | 16 (20.1%)                           | 0.65 (0.25–1.64)                          |
| Gestational diabetes   | 8 (11.1%)                                      | 10 (13.0%)                           | 1.01 (0.34–2.99)                          |
| Antepartum hemorrhage/abruption                                  | 7 (9.7%)                                       | 7 (9.1%)                             | 0.92 (0.28–3.0)                           |

Data expressed as number (percentage) unless otherwise indicated.

CI, confidence interval; NA, not applicable; SGA, small for gestational age; HELLP, haemolysis, elevated liver enzymes, low platelets.

<sup>a</sup> Multiple logistic regression adjusted for inclusion criteria, recruitment center, final thrombophilia status, and number of previous preeclampsia/SGA events (1 vs  $\geq 2$  events); <sup>b</sup> Event rate too low for formal testing.Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.**TABLE 3**  
**Delivery outcomes**

|                                      | Standard high-risk care and enoxaparin, n = 72 | Standard high-risk care only, n = 77 | Adjusted odds ratio <sup>a</sup> (95% CI) or Pvalue |
|--------------------------------------|--|--------------------------------------|---|
| Preterm birth $\leq 36^{+6}$ wk      | 16 (22.2%)                                     | 19 (24.7%)                           | 0.76 (0.33–1.73)                                    |
| Preterm birth $\leq 33^{+6}$ wk      | 7 (9.7%)                                       | 10 (13.0%)                           | 0.55 (0.17–1.75)                                    |
| Induction of labor                   | 15 (20.8%)                                     | 17 (22.1%)                           | 0.76 (0.33–1.73)                                    |
| Cesarean delivery                    | 39 (54.1%)                                     | 40 (52.0%)                           | 1.35 (0.68–2.71)                                    |
| Prelabor cesarean delivery           | 31 (43.1%)                                     | 32 (41.6%)                           | 1.34 (0.66–2.72)                                    |
| Major postpartum hemorrhage >1000 mL | 4 (5.6%)                                       | 7 (9.1%)                             | <sup>b</sup>  |
| Postpartum hemorrhage $\geq 500$ mL  | 27 (37.5%)                                     | 30 (39.0%)                           | 0.79 (0.38–1.63)                                    |
| Fetal loss >16 wk                    | 2 (2.8%) <sup>c</sup>                          | 3 (3.9%) <sup>d</sup>                | <sup>b</sup>  |
| Gestational age at delivery, d       | 268 (118–292)                                  | 269 (147–292)                        | .68   |
| Birthweight, g                       | 2920 (22–4582)                                 | 3150 (385–4120)                      | .70   |
| Birthweight percentile               | 17.1 (0–99)                                    | 22.9 (0–94)                          | .93   |

Data expressed as median (minimum–maximum values) or number (percentage) unless otherwise indicated.

CI, confidence interval.

<sup>a</sup> Multiple logistic regression adjusted for inclusion criteria, recruitment center, final thrombophilia status, and number of previous preeclampsia/small-for-gestational-age events (1 vs  $\geq 2$  events); <sup>b</sup> Event rate too low for formal testing; <sup>c</sup> Includes termination of pregnancy at 17 wk gestation for severe early-onset intrauterine growth restriction and stillbirth at 25<sup>+</sup> wk with severe early-onset intrauterine growth restriction; <sup>d</sup> Includes stillbirth at 24<sup>+</sup> wk with abruption, stillbirth at 20<sup>+</sup> wk after extreme preterm birth, and stillbirth at 38<sup>+</sup> wk with severe preeclampsia, small-for-gestational-age <10th percentile, and abruption.Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.

TABLE 4

## Neonatal outcomes (for live-born only)

|   | Standard high-risk care and enoxaparin, n = 70 | Standard high-risk care only, n = 74 | Adjusted odds ratio <sup>a</sup> (95% CI) or Pvalue |
|---|--|--------------------------------------|---|
| Neonatal death                              | 0  | 0                                    | NA  |
| NICU admission                              | 14 (20%)                                       | 14 (18.9%)                           | 0.93 (0.37–2.37)                                    |
| Duration of NICU admission, d               | 13.5 (1–116)                                   | 23.5 (5–126)                         | .37   |
| Composite outcome severe neonatal morbidity | 2 (2.9%)                                       | 4 (5.4%)                             | <sup>b</sup>  |

Data expressed as median (minimum–maximum values) or number (percentage) unless otherwise indicated.

CI, confidence interval; NA, not applicable; NICU, neonatal intensive care unit.

<sup>a</sup> Multiple logistic regression adjusted for inclusion criteria, recruitment center, final thrombophilia status, and number of previous PET/small-for-gestational-age events (1 vs  $\geq 2$  events); <sup>b</sup> Event rate too low for formal testing.

Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.

composite primary outcome. Our inclusion criteria resulted in a higher proportion of participants with a history of a SGA compared to other trials. This is likely to explain the relatively low rates of recurrence of preeclampsia compared to SGA. Subgroup analyses (although not sufficiently powered to this end) did not indicate any effect of LMWH is limited to preeclampsia and not SGA.

The result of our biomarker studies further supports the premise that LMWH does not have a therapeutic effect for preeclampsia only. An important marker of preeclampsia is the elevated placental release of the antiangiogenic factors sFlt-1 and sEng. They spread throughout the maternal circulation and cause widespread endothelial dysfunction, and blood vessel damage causing hypertension and multiorgan injury. We have shown there were no changes in serum sFlt-1 and sEng concentrations

among women treated with enoxaparin, compared to standard high-risk care only. Importantly, we did observe an expected rise in sFlt-1 and sEng among those who subsequently developed preeclampsia, which was not mitigated by the use of enoxaparin. We also found enoxaparin was unable to decrease circulating levels of ET-1 (a potent endothelial-derived vasoconstrictor known to be increased in preeclampsia) among those who subsequently developed preeclampsia (and SGA).

It may be argued that the estimation of the size of effect of LMWH used in our sample size calculation was unrealistic. However, the estimated recurrence rate for the standard high-risk care only group (25%) was very similar to the overall recurrence rate (23.5%) and indeed the same as the rate of recurrence for those treated with additional LMWH (25.0%). Our estimate for a reduction in

risk of recurrence to 7% was optimistic but based on the most applicable data set published at that time, which reported a similar composite primary outcome of 5.5%.<sup>22</sup> This was an open-label trial with no placebo arm. Participants, clinicians, and investigators were aware of the assigned treatment group. We originally planned a placebo-controlled trial, however, we were unable to commercially source suitable placebo syringes and were unable to secure sterile manufacture and preparation of syringes that would meet international regulatory requirements. No previous trials of LMWH for the prevention of placenta-mediated pregnancy complications used placebo injections as a comparator and, even in nonpregnant trials of LMWH, placebo has only been used in industry-sponsored trials. We therefore elected to proceed with an open-label trial with no placebo injection. Our

TABLE 5

## History subgroup analysis of primary outcome

|  | Standard high-risk care and enoxaparin | Standard high-risk care only | Unadjusted odds ratio <sup>a</sup> (95% CI) |
|--|--|------------------------------|---|
| Preeclampsia $\pm$ SGA <sup>b</sup> n = 63                                 | 8/30 (26.7%)                           | 6/33 (18.2%)                 | 1.64 (0.49–5.43)                            |
| SGA infant $\leq 3$ rd percentile with no preeclampsia <sup>b</sup> n = 65 | 9/34 (26.5%)                           | 8/31 (25.8%)                 | 1.04 (0.34–3.13)                            |
| SGA infant $< 10$ th percentile with no preeclampsia <sup>b</sup> n = 79   | 9/38 (23.7%)                           | 9/41 (22.0%)                 | 1.10 (0.39–3.16)                            |

Data expressed as number with primary outcome/total (percentage) unless otherwise indicated.

CI, confidence interval; SGA, small for gestational age.


<sup>a</sup> Unadjusted crude odds ratio used as numbers too small for appropriate adjustment; <sup>b</sup> In last ongoing pregnancy reaching  $> 12$  wk.

Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.



FIGURE 2

Serum biomarker levels in women treated with enoxaparin or standard high-risk care only taken at baseline (recruitment), 20 weeks, and 30 weeks



Serum levels of **A**, soluble fms-like tyrosine kinase (sFlt)-1; **B**, soluble endoglin (sEng); **C**, placental growth factor (PlGF); **D**, soluble vascular cell adhesion molecule (VCAM)-1; and **E**, endothelin (ET)-1 taken at baseline (recruitment), 20 weeks, and 30 weeks were similar for those receiving enoxaparin (treated) and those receiving standard high-risk care only (nontreated). Serum biomarker levels according to treatment group. \* $P < 0.05$ ; black closed circles indicate standard high risk care only; blue closed circles indicate enoxaparin treatment.

Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. Am J Obstet Gynecol 2017.


trial had similar issues to other trials with a prolonged recruitment phase. Following a feasibility assessment at the lead trial site it was estimated that with a 30% recruitment rate we would complete recruitment within 2 years. However, a large number of women were not referred until  $>16^{+0}$  weeks' gestation and so were not eligible for the trial. Trial duration was extended and additional recruiting sites were identified to ensure the trial was completed to the planned sample size.

Our results conflict with those seen in a published pooled summary-based, study-level meta-analysis including 6 trials and 848 pregnant women. A relative risk reduction of 0.52 (95% CI, 0.32–0.86; 18.7% vs 42.9%) was demonstrated for a primary composite outcome (preeclampsia, SGA  $<10$ th percentile, placental abruption, or pregnancy loss  $>20$  weeks) with similar risk reductions for a number of secondary outcomes including preeclampsia, SGA, and preterm birth.<sup>29</sup> However, this meta-analysis showed high heterogeneity ( $I^2$  69%) and the included trials of higher quality suggested no treatment effect. Individual patient data analysis including these same trials and 3 additional trials showed significant heterogeneity between single-center and multicenter trials with no effect seen on a primary composite outcome (early-onset or severe preeclampsia, SGA  $<5$ th percentile, placental abruption, or pregnancy loss  $>20$  weeks) for multicenter trials (18% LMWH treatment vs 18% no LMWH treatment) but a significant reduction in single-center trials (8% vs 27% respectively, absolute difference  $-18.7\%$ ; 95% CI,  $-21.6$  to  $-15.7$ ;  $P < .0001$ ).<sup>30</sup>

The multicenter EPPI trial was designed to be as specific to preeclampsia and SGA as possible. Thrombophilia status was accounted for but did not limit the trial population. Women in both groups consistently received standard high-risk obstetric care. Our trial findings are consistent with the 2 most recently published multicenter trials of LMWH. Both the HAPPY<sup>25</sup> and HEPEPE<sup>26</sup> trials included women at high risk of placental-mediated

FIGURE 3

**Serum biomarker levels in women who developed preeclampsia (PE) compared to those who did not taken at baseline (recruitment), 20 weeks, and 30 weeks**



There was significant increase in levels of **A**, soluble fms-like tyrosine kinase (sFlt)-1; **B**, soluble endoglin (sEng); **C**, placental growth factor (PlGF); and **D**, soluble vascular cell adhesion molecule (VCAM)-1 at 30 weeks' gestation observed in women who developed PE compared to those who did not. **E**, Significant increase in endothelin (ET)-1 levels was observed in women who developed PE across all time points examined at baseline (recruitment) and at 20 and 30 weeks' gestation. Serum biomarker levels according to preeclampsia status. \* $P < 0.05$ ; \*\* $P < 0.01$ ; black closed circles indicate no preeclampsia; blue closed circles indicate developed preeclampsia.

Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.

pregnancy complications, with or without thrombophilia, and reported no difference in rates of preeclampsia or SGA associated with LMWH use. It seems likely these 3 trials are more representative of a general at-risk population. Data from the EPPI trial will make a significant contribution to future meta-analyses.

## Acknowledgment

Participating centers: National Women's Health, Auckland City Hospital, Auckland, New Zealand; Department of Maternal-Fetal Medicine, Royal Women's Hospital, Melbourne, Australia; Department of Perinatal Medicine, Mercy Hospital for Women, Melbourne, Australia; Department of Obstetrics, Academic Medical Center, Amsterdam, The Netherlands; and Department of Maternal Fetal Medicine, Sunshine Hospital, Melbourne, Australia.

EPPI Trial Investigator Group. University of Auckland and Auckland City Hospital: K. Groom, L. McCowan, L. Mackay, A. Lee, P. Stone, L. Chamley, C. McLintock. Royal Women's Hospital and University of Melbourne: J. Said, S. Kane. Mercy Hospital for Women and University of Melbourne: S. Walker, S. Tong, N. Hannan. Sunshine Hospital and University of Melbourne: J. Said. Academic Medical Center: T. van Mens, W. Ganzevoort, S. Middeldorp.

We thank all the women who participated in the EPPI Trial, the research staff (K. Anderson, L. Wilkinson, J. Callender, and L. Harris, University of Auckland; D. Maxwell and M. Poth, Royal Women's Hospital; G. Pell, University of Melbourne, Mercy Hospital for Women; and L. Lynch, Maternal Fetal Medicine Unit, Sunshine Hospital), and the data monitoring committee (A. Stewart, University of Auckland; S. Belgrave, Auckland City Hospital and North Shore Hospital; J. Rowan, Auckland City Hospital).

This trial was presented to, and peer reviewed by, the Perinatal Society of Australia and New Zealand Interdisciplinary Maternal Perinatal Australasian Collaborative Trials Clinical Trials (IMPACT) Network for Mothers' and Babies' Health.

## References

- McCowan LM, Harding JE, Stewart AW. Customized birthweight centiles predict SGA pregnancies with perinatal morbidity. *BJOG* 2005;112:1026-33.
- Vintzileos AM, Ananth CV. First trimester prediction of ischemic placental disease. *Semin Perinatol* 2014;38:159-66.
- McCowan LM, Thompson JM, Taylor RS, et al. Clinical prediction in early pregnancy of infants small for gestational age by customized birthweight centiles: findings from a healthy nulliparous cohort. *PLoS One* 2013;8:e70917.

4. Duffett L, Rodger M. LMWH to prevent placenta-mediated pregnancy complications: an update. *Br J Haematol* 2015;168:619-38.
5. Hnat MD, Sibai BM, Caritis S, et al. Perinatal outcome in women with recurrent preeclampsia compared with women who develop preeclampsia as nulliparas. *Am J Obstet Gynecol* 2002;186:422-6.
6. Makkonen N, Heinonen S, Kirkinen P. Obstetric prognosis in second pregnancy after preeclampsia in first pregnancy. *Hypertens Pregnancy* 2000;19:173-81.
7. Sibai BM, el-Nazer A, Gonzalez-Ruiz A. Severe preeclampsia-eclampsia in young primigravid women: subsequent pregnancy outcome and remote prognosis. *Am J Obstet Gynecol* 1986;155:1011-6.
8. van Rijn BB, Hoeks LB, Bots ML, Franx A, Bruinse HW. Outcomes of subsequent pregnancy after first pregnancy with early-onset preeclampsia. *Am J Obstet Gynecol* 2006;195:723-8.
9. Askie LM, Duley L, Henderson-Smart DJ, Stewart LA; PARIS Collaborative Group. Antiplatelet agents for prevention of pre-eclampsia: a meta-analysis of individual patient data. *Lancet* 2007;369:1791-8.
10. Hofmeyr GJ, Lawrie TA, Atallah AN, Duley L, Torloni MR. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *Cochrane Database Syst Rev* 2014;6:CD001059.
11. Duley L, Henderson-Smart DJ, Meher S, King JF. Antiplatelet agents for preventing preeclampsia and its complications. *Cochrane Database Syst Rev* 2007;2:CD004659.
12. Sibai B, Dekker G, Kupferminc M. Preeclampsia. *Lancet* 2005;365:785-99.
13. National Institute for Health and Clinical Excellence. Hypertension in pregnancy: diagnosis and management. Clinical guideline [CG107]. Available at: <https://www.nice.org.uk/guidance/cg107/chapter/1-Guidance#reducing-the-risk-of-hypertensive-disorders-in-pregnancy>. Accessed January 25th, 2010.
14. Lowe SA, Bowyer L, Lust K, et al. The SOMANZ guidelines for the management of hypertensive disorders of pregnancy. Available at: <https://somanz.org/documents/HTPregnancyGuidelineJuly2014.pdf>. Accessed May 20, 2014.
15. Quenby S, Mountfield S, Cartwright JE, Whitley GS, Vince G. Effects of low-molecular-weight and unfractionated heparin on trophoblast function. *Obstet Gynecol* 2004;104:354-61.
16. Bose P, Black S, Kadyrov M, et al. Heparin and aspirin attenuate placental apoptosis in vitro: implications for early pregnancy failure. *Am J Obstet Gynecol* 2005;192:23-30.
17. Bose P, Black S, Kadyrov M, et al. Adverse effects of lupus anticoagulant positive blood sera on placental viability can be prevented by heparin in vitro. *Am J Obstet Gynecol* 2004;191:2125-31.
18. North RA, Ferrier C, Gamble G, Fairley KF, Kincaid-Smith P. Prevention of preeclampsia with heparin and antiplatelet drugs in women with renal disease. *Aust N Z J Obstet Gynaecol* 1995;35:357-62.
19. Mello G, Parretti E, Fatini C, et al. Low-molecular-weight heparin lowers the recurrence rate of preeclampsia and restores the physiological vascular changes in angiotensin-converting enzyme DD women. *Hypertension* 2005;45:86-91.
20. Kupferminc MJ, Fait G, Many A, et al. Low-molecular-weight heparin for the prevention of obstetric complications in women with thrombophilias. *Hypertens Pregnancy* 2001;20:35-44.
21. Riyazi N, Leeda M, de Vries JI, Huijgens PC, van Geijn HP, Dekker GA. Low-molecular-weight heparin combined with aspirin in pregnant women with thrombophilia and a history of preeclampsia or fetal growth restriction: a preliminary study. *Eur J Obstet Gynecol Reprod Biol* 1998;80:49-54.
22. Rey E, Garneau P, David M, et al. Dalteparin for the prevention of recurrence of placental-mediated complications of pregnancy in women without thrombophilia: a pilot randomized controlled trial. *J Thromb Haemost* 2009;7:58-64.
23. de Vries JI, van Pampus MG, Hague WM, Bezemer PD, Joosten JH; FRUIT Investigators. Low-molecular-weight heparin added to aspirin in the prevention of recurrent early-onset preeclampsia in women with inheritable thrombophilia: the FRUIT-RCT. *J Thromb Haemost* 2012;10:64-72.
24. Rodger MA, Hague WM, Kingdom J, et al. Antepartum dalteparin versus no antepartum dalteparin for the prevention of pregnancy complications in pregnant women with thrombophilia (TIPPS): a multinational open-label randomized trial. *Lancet* 2014;384:1673-83.
25. Martinelli I, Ruggerenti P, Cetin I, et al. Heparin in pregnant women with previous placenta-mediated pregnancy complications: a prospective, randomized, multicenter, controlled clinical trial. *Blood* 2012;119:3269-75.
26. Haddad B, Winer N, Chitrit Y, et al. Enoxaparin and aspirin compared with aspirin alone to prevent placenta-mediated pregnancy complications: a randomized controlled trial. *Obstet Gynecol* 2016;128:1053-63.
27. Visser J, Ulander VM, Helmerhorst FM, et al. Thromboprophylaxis for recurrent miscarriage in women with or without thrombophilia. HEBE-NOX: a randomized multicenter trial. *Thromb Haemost* 2011;105:295-301.
28. Kaandorp SP, Goddijn M, van der Post JA, et al. Aspirin plus heparin or aspirin alone in women with recurrent miscarriage. *N Engl J Med* 2010;362:1586-96.
29. Rodger MA, Carrier M, Le Gal G, et al. Meta-analysis of low-molecular-weight heparin to prevent recurrent placenta-mediated pregnancy complications. *Blood* 2014;123:822-8.
30. Rodger MA, Gris JC, de Vries JI, et al. Low-molecular-weight heparin and recurrent placenta-mediated pregnancy complications: a meta-analysis of individual patient data from randomized controlled trials. *Lancet* 2016;388:2629-41.
31. Levine RJ, Maynard SE, Qian C, et al. Circulating angiogenic factors and the risk of preeclampsia. *N Engl J Med* 2004;350:672-83.
32. Maynard SE, Min JY, Merchan J, et al. Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J Clin Invest* 2003;111:649-58.
33. Venkatesha S, Toporsian M, Lam C, et al. Soluble endoglin contributes to the pathogenesis of preeclampsia. *Nat Med* 2006;12:642-9.
34. Schiff E, Ben-Baruch G, Peleg E, et al. Immunoreactive circulating endothelin-1 in normal and hypertensive pregnancies. *Am J Obstet Gynecol* 1992;166:624-8.
35. Sanofi-Aventis. Clexane and Clexane Forte. Medsafe. Available at: <http://www.medsafe.govt.nz/profs/datasheet/c/clexaneinj.pdf>. Accessed July 29th, 2016.
36. Samama MM, Cohen AT, Darmon JY, et al. A comparison of enoxaparin with placebo for the prevention of venous thromboembolism in acutely ill medical patients. Prophylaxis in Medical Patients with Enoxaparin Study Group. *N Engl J Med* 1999;341:793-800.
37. Salomon LJ, Alfirevic Z, Berghella V, et al. Practice guidelines for performance of the routine mid-trimester fetal ultrasound scan. *Ultrasound Obstet Gynecol* 2011;37:116-26.
38. Bhide A, Acharya G, Bilardo CM, et al. ISUOG practice guidelines: use of Doppler ultrasonography in obstetrics. *Ultrasound Obstet Gynecol* 2013;41:233-9.
39. Tranquilli AL, Dekker G, Magee L, et al. The classification, diagnosis and management of the hypertensive disorders of pregnancy: a revised statement from the ISSHP. *Pregnancy Hypertens* 2014;4:97-104.
40. Anderson NH, Sadler LC, Stewart AW, McCowan LM. Maternal and pathological pregnancy characteristics in customized birthweight centiles and identification of at-risk small-for-gestational-age infants: a retrospective cohort study. *BJOG* 2012;119:848-56.
41. Abbassi-Ghanavati M, Greer LG, Cunningham FG. Pregnancy and laboratory studies: a reference table for clinicians. *Obstet Gynecol* 2009;114:1326-31.
42. Gris JC, Chauleur C, Faillie JL, et al. Enoxaparin for the secondary prevention of placental vascular complications in women with abruptio placentae. The pilot randomized controlled NOH-AP trial. *Thromb Haemost* 2010;104:771-9.

## Author and article information

From the Department of Obstetrics and Gynecology (Drs Groom, McCowan, Chamley, and Stone, and Ms Mackay) and Section of Epidemiology and Biostatistics (Dr Lee), University of Auckland, and National Women's Health, Auckland City Hospital (Drs Groom, McCowan, and McLintock), Auckland, New Zealand; Department of Obstetrics and Gynecology, University of Melbourne (Drs Said, Kane, and Walker), Department of Maternal-Fetal

Medicine, Sunshine Hospital (Dr Said), Pregnancy Research Center, Department of Maternal-Fetal Medicine, Royal Women's Hospital (Dr Kane), Mercy Hospital for Women (Dr Walker), and Translational Obstetrics Group, Department of Obstetrics and Gynecology, Mercy Hospital for Women, University of Melbourne (Drs Hannan and Tong), Melbourne, Australia; and Department of Vascular Medicine, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands (Dr van Mens).

Received Nov. 14, 2016; revised Jan. 5, 2017; accepted Jan. 13, 2017.

Supported by University of Auckland Faculty Research Development Fund (3625687/9803 and 3624720/9803); A+ Trust, Auckland District Health Board Charitable Trust (A+2011056); Australasian Society of Thrombosis and Hemostasis Scientific and Education Trust (3708366/9103); and the University of Auckland School of Medicine Performance Based Research Fund (71603/9450). The funders had no role in study design;

in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the article for publication.


The authors report no conflict of interest.

Presented at the 37th annual meeting of the Society for Maternal-Fetal Medicine, Las Vegas, NV, Jan. 23-28, 2017.

Corresponding author: Katie M. Groom, MBBS, PhD, FRANZCOG, CMFM. [k.groom@auckland.ac.nz](mailto:k.groom@auckland.ac.nz)

## SUPPLEMENTARY FIGURE 1

Serum biomarker levels in women who developed preeclampsia taken at baseline (recruitment), 20 weeks, and 30 weeks treated with enoxaparin or standard high-risk care only




Levels of **A**, soluble fms-like tyrosine kinase (sFlt)-1; **B**, soluble endoglin (sEng); **C**, placental growth factor (PlGF); **D**, soluble vascular cell adhesion molecule (VCAM)-1; and **E**, endothelin (ET)-1 were not significantly changed by treatment with enoxaparin (treated) among women who developed preeclampsia compared to those receiving standard high-risk care only (nontreated). Black closed circles indicate standard high-risk care; blue closed circles indicate enoxaparin.

Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol* 2017.



## SUPPLEMENTARY FIGURE 2

Serum biomarker levels in women who developed small for gestational age (SGA) <5th percentile compared to those with infants with birthweight  $\geq$ 5th percentile taken at baseline (recruitment), 20 weeks, and 30 weeks



Levels of **A**, soluble fms-like tyrosine kinase (sFlt)-1; **B**, soluble endoglin (sEng); **C**, placental growth factor (PlGF); and **D**, soluble vascular cell adhesion molecule (VCAM)-1 were similar for those with SGA <5th percentile and those with infant birthweight  $\geq$ 5th percentile at all time points. **E**, There was significant increase in endothelin (ET)-1 levels at baseline in serum from women who later developed SGA pregnancy. \*P<0.05; black closed circles indicate non-small for gestational age; blue closed circles indicate small for gestational age.

Groom et al. Enoxaparin for prevention of preeclampsia and intrauterine growth restriction. Am J Obstet Gynecol 2017.